

Full Body Immersive Virtual Reality System with Motion Recognition Camera targeting the treatment of Spider Phobia

Jacob Kritikos^{*1}, Stavroula Pouloupoulou², Chara Zoitaki³, Marilina Douloudi^{4*}, Prof. Dimitris Koutsouris^{*1}

Abstract. Exposure Therapy (ET) is one of the most widely-used methods for treating Specific Phobias, and over the past few years, Virtual Reality (VR) has contributed significantly in this field, since the birth of what we call ‘Virtual Reality ET’ (VRET). However, VR systems used in VRET so far do not fully integrate ET characteristics; the reason behind this is that they do not provide sufficient, or occasionally not any at all, interaction with the feared stimulus, which is a key factor for full ET implementation. Objective: The aim of this paper is to propose a way to include natural interaction between the patient and the system both in the treatment procedure and the VR system. Method: We propose an addition to current session protocols for mental health professionals through which they can apply ET in full extent with the use of Motion Tracking Sensors. Specifically, we added a Motion Recognition Camera, which tracks the patient’s movements and places their physical body within the virtual environment, increasing their feeling of presence and making the system more immersive. Therefore, clinicians can assign interactive tasks for their patients to practice within a controlled virtual environment. Results: We present the results of a trial run by a group of psychiatry professionals, who provided feedback on its utility and efficiency. Clinical Impact: Patients can truly confront their phobias and acquire the necessary skills to overcome them.

Keywords: Virtual Reality, Cognitive Behavioral Treatments, Exposure Therapy, Anxiety Disorders, Specific Phobias, Motion Tracking Sensor, Motion Recognition Camera, Presence, Clinical Treatment, Immersion

* Corresponding authors. Address: Iroon Polytechniou 9, N.T.U.A (Zografou campus), Zografou, 15780 Athens, Greece. Tel.: +30 2107722430. E-mail addresses: jkrikos@biomed.ntua.gr (J. Kritikos), dkoutsou@biomed.ntua.gr (D. Koutsouris).

¹ Biomedical Engineering Laboratory, School of Electrical and Computer Engineering, National Technical University of Athens, Athens, Greece

² Department of Digital Systems, University of Piraeus, Piraeus, Greece

³ School of Electrical and Computer Engineering, National Technical University of Athens, Athens, Greece

⁴ Biology dept. National Kapodistrian University of Athens, Greece

1 Introduction

Anxiety disorders share features of excessive worry and fear that cause behavioral disturbances to the individuals suffering. Approximately 1 in 5 adults will manifest some form of Anxiety Disorder at some point in their lives with rates being twice as high for minors (American Psychiatric Association, 2013; Bandelow & Michaelis, 2015; *National Institute of Mental Health - Anxiety Disorder*, n.d.; *National Institute of Mental Health - Mental Illness*, n.d.). Specific Phobias are a type of an anxiety disorder defined as the persistent fear of a stimulus, potentially an object or a situation, which renders the person unable to demonstrate self-control. Specific Phobias along with Agoraphobia and Social phobia are one of the most common types of phobias, which in fact are considered as sub-types of a broader category, that of anxiety disorders, as stated by the Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-V). According to the American Psychiatric Association (American Psychiatric Association, 2013, p. 199), the lifetime prevalence of Specific Phobias is approximately 7%-9% in the US and 6% in Europe, and can reach up to 12,5% (Kessler et al., 2005). Although a lot of research has been conducted in this field over the past few years, only a minority (8%) of people reported with Specific Phobias have received any treatment (Öst, 1989; Stinson et al., 2007).

The most well-known treatment of various cognitive and behavioral disorders is Cognitive Behavioral Therapy (CBT) (Mayo-Wilson & Montgomery, 2013; Scott & Beck, 2008), since it appears as the preferred method by the majority of mental health professionals to treat phobias; this may be due to the fact that a number of scientists claim CBT has no possible side-effect, unlike medication (American Psychiatric Association, 2013) (“Treatment - (ADAA) Anxiety and Depression Association of America,” n.d.). The basic principles of CBT focus on identifying, understanding and altering patients’ thought and behavioral patterns. Patients are actively involved in their own recovery, which offers them a stronger sense of control; additionally, through therapy sessions, they obtain useful skills they ought to practice repeatedly in order to present the desirable progress (Otte, 2011).

The applications of CBT, along with its different forms, have been highly efficient so far; by evaluating previous studies, we can deduce that an average of 50% of patients managed to decrease their phobia-related anxiety (Kerns, Read, Klugman, & Kendall, 2013; Krebs et al., 2017; Riddle-Walker et al., 2016; Rozentel et al., 2017; Warwick et al., 2017; Wittchen, Mühlhig, & Beesdo, 2003). Nonetheless, certain studies reviewing CBT results (Peñate et al., 2017; Twomey, O’Reilly, & Meyer, 2017) indicate that it is still unclear whether the phobia returns after a short period of time or not. For instance, a recent research (Ollendick, Öst, Ryan, Capriola, & Reuterskiöld, 2017) claims that the phobia does not return at least amid the first semester. Yet, some scientists disclose that CBT’s benefits do not necessarily last long; specifically, another study (Shiban, Schelhorn, Pauli, & Mühlberger, 2015) mentions that results lasted only for the first 20 days.

Exposure Therapy (ET) is a form of CBT treatment (Treanor & Craske, 2015) developed to help people confront their fears in a safe environment. There are several variations of ETs, such as ‘In VIVO ET’, ‘Virtual Reality ET’ (VRET) (Rothbaum, Rizzo, McDaniel, & Zanol, 2015), ‘Imaginary ET’ and ‘Interoceptive ET’ (Rothbaum, Rizzo, McDaniel, & Zanol, 2015; (Society of Clinical Psychology, 2013) but all of them follow the same

basic procedure (Cardoş, David, & David, 2017; Rizzo, Hartholt, Grimani, Leeds, & Liewer, 2014; Waters et al., 2014), which usually consists of the following steps:

- (Step 1): The Cognitive sessions, where the patient is informed about their condition and their unreasonable thought patterns; next, the patient discusses any existent distressing thoughts and possible dreaded scenarios with the experts.
- (Step 2): The exposure sessions, where the patient is exposed to assorted phobic situations with graded difficulty through In VIVO ET or VRET. This step is repeated until the patient's anxiety levels decrease.
- (Step 3): The follow-up session appointed after a significant period of time in which clinicians evaluate the overall decrease in anxiety levels.

During In VIVO ET, patients are called to confront the feared stimulus in real life. However, simulating every possible scenario in real life is expensive and time-consuming; for instance, in case a patient was suffering from aviophobia, they would have to buy an airplane ticket and go through the entire process accompanied by their clinician, who charges by the hour (Bun, Gorski, Grajewski, Wichniarek, & Zawadzki, 2016; Yiasemidou et al., 2017). Also, in some cases, it could lead to undesired results since displeasing occurrences may further terrorize patients (e.g., turbulence during the flight in the early stages of ET (Blakey & Abramowitz, 2016)). In order to overcome those technical issues, VRET is starting to replace In VIVO ET.

In VRET, patients are exposed to virtual, life-like, anxiety-provoking environments instead of real stimuli (Carlin, Hoffman, & Weghorst, 1997; Emmelkamp et al., 2002; Krijn et al., 2004). Within them, they can practice tasks assigned by their clinician in a controllable, and therefore safe, virtual environment designed to appropriately stimulate their Specific Phobia, in the clinician's office. Due to that, patients feel more comfortable and confident; thus, they are more willing to confront situations that cause them discomfort and explore alternative ways of responding. Also, even though their actions occur in the virtual world, the knowledge acquired transfers to the real world (Daniel Freeman et al., 2018). Lastly, over the last few years, studies have found VRET to be as effective as In VIVO (Krijn et al., 2004) in terms of triggering anxiety. Therefore, treatment through VR systems could become a low-cost method of providing effective interventions at scale.

However, looking at current applications, we notice that all VRET methods applied so far have not sufficiently integrated ET's two primary characteristics: [a] patients have to come in contact with the feared stimulus actively, and [b] patients have to master how to confront and respond to stimuli without fear or anxiety. Current VRET applications are only able to integrate the first characteristic, i.e. to trigger fear through VR videos; however, they have not yet been able to assist patients in interacting with the stimulus. At best, the only interaction that is currently available is via Hand Controllers through which complex tasks and exercises, such as touching and moving objects with their joints (e.g. their elbows or feet) cannot be completed, due to the nature of Hand Controllers' design that only provides hand interaction; therefore, certain scenarios cannot be simulated. Overall, VRET has managed to stimulate patients' anxiety through a virtual stimulus; yet, patients usually remain passive or interact poorly, which doesn't improve immersion, and consequently, the patient's feeling of presence in the virtual environment is decreased.

Hence, in order for VRET to reach its full potential and implement ET as a whole, life-like interaction must be added to existing systems (Krijn et al., 2004; Morina, Brinkman, Hartanto, & Emmelkamp, 2014; Price & Anderson, 2007). Therefore, in this study, we present and propose a fully immersive VR System by adding a new tool, the Motion Recognition Camera (MRC), on top of the VR technology. The MRC tracks patients' movements, places their physical body within the virtual environment and gives them the impression that they are moving and interacting in full extent with that environment, as they would in the real world. This allows patients to practice tasks whilst in the virtual environment. By combining these technologies, we increase the user's immersion and presence. Additionally, we propose that Step 2 be separated into two different cycles regarding the Exposure sessions: [a] Step 2-a in which Exposure Therapy is applied as it currently is; patients come in contact with the phobic stimulus for the first time in a controlled environment and learn how to stay close to the stimulus without losing their composure, and [b] Step 2-b in which patients practice possible coping mechanisms, that differ from the ones originated from physical environment, like concepts of avoidance and fear anticipation (Rudaz, Ledermann, Margraf, Becker, & Craske, 2017) by directly interacting with the phobic stimulus, and by learning to control their reaction and regulate their anxiety response. In this study, we identify Step 2-b as Action Therapy for we want to emphasize the importance of confrontation, which has to become a separate and independent part of the sessions.

So, our hypothesis is whether the proposed system can provide appropriate interaction and confrontation with the phobic stimulus, which could improve current treatments in the future. We examined this hypothesis through a trial on a psychiatric clinic's medical staff. After completing the study, the clinicians gave us their professional feedback on their experience with the proposed system as well as suggestions for further improvement.



Figure 1: The MRC which is in front of the participant recognize the joints and place those inside the virtual world.

2 Setup & Materials

2.1 Hardware

For the execution of our trials we used the following equipment: [a] VR Goggles (VRG): "Gear VR" by Samsung, [b] a Mobile Phone: the "Galaxy S7" by Samsung, [c] a Motion Recognition Camera (MRC): the "Astra S" by ORBBEC; the specifications of the MRC are: Range: 0.6-8.0m (Optimal 0.6-5.0m), Depth Image Size: 640*480-30fps, RGB Image Size: 1280*960-30fps, Field of View: 60° horiz. x 49.5° vert. (73° diagonal). The MRC tracks the patient's movements, places their physical body in the virtual environment and gives them the impression that they are moving and interacting with their whole body in that environment in real time, as they would in the real world. This allows the patient to practice tasks assigned by their clinician in the virtual environment; [d] a Windows Desktop Computer.

2.2 Software

We created a C++ program and used the Orbbec Astra SDK, along with the Bluetooth Windows SDK, to assist us in body recognition and its dispatching from the Desktop Computer, which operates the MCR, to the Gear VR. Android Studio was used to create the Bluetooth receiver program; Unity was used for creating the virtual environment depicted by the VRG; Blender 3D Computer Graphics was used for creating 3D objects, animated visual effects and materials, and Adobe Photoshop for designing images for




Figure 2: The participant can see his skeleton inside the virtual room. (*top*) Here is the participant's right arm; (*bottom*) Here is the participant's left arm.

the materials. Moreover, regarding the body Tracking Software, the Astra S MRC is designed for skeleton tracking (Figure 1); consequently, it can recognize the entire body and movements of its limbs. The program we have written for the MRC tracks the joints (Figure 2); then, the user's skeleton is represented in the virtual room as a set of spheres and lines (the spheres represent the joints and the lines represent the bones connecting joints).

2.3 Setup

The outlined area is about 5 m² and will be referred to as the 'Action Area'. In the Action Area, the user wears the VRG and is thus able to observe the virtual environment. The user does not need to wear or hold any other equipment than the VRG, in order to move and interact with objects. The MRC is placed on a table at a 1 m distance from the Action Area, so as to record the user's movements and transfer them to the virtual environment.

3 Method

The proposed system was tested by twenty members of the Psychiatry Department of the Oncological Hospital of Kifisia staff (doctors, professors, nursing staff and the hospital's general manager), in Athens, Greece. All of the participants were familiar with the concept of using visual cues before (pictures, videos, standalone VR) to stimulate fear to phobic patients. The aim of the session was for all participants to use the proposed system and answer a small questionnaire on their experience with it.

Q1: Do you believe that Exposure Therapy could benefit from the addition of interaction?

_____	_____	_____	_____	_____
NO	MAYBE-NO	MAYBE	MAYBE-YES	YES

Q2: Do you think that interaction with the Motion Recognition Camera in the Virtual Environment can give real life experience?

_____	_____	_____	_____	_____
NO	MAYBE-NO	MAYBE	MAYBE-YES	YES

Q3: Do you believe that coping mechanisms learnt in virtual situations and environments can be transferred to real-life situations?

_____	_____	_____	_____	_____
NO	MAYBE-NO	MAYBE	MAYBE-YES	YES

Furthermore, we present an implementation of Action Therapy. Spider phobia used as a case study, which consists of one session with two levels of difficulty. During this session, two tasks are assigned to participants; their aim is to assist participants in mastering new coping mechanisms, so as to confront their phobia; to habituate with the phobic environment and eventually interact with the stimulus through their own pace and control.

Level 01: Step 1: The participants enter an empty 5 m² area with gradients, while the MRC is located at a 1m distance. This is the area where the patient can walk, move and react to the stimuli. **Step 2:** The participant wears the VRG and enters the virtual environment (Figure 3). **Step 3:** In the VR room, the patient can observe their virtual arms, as well as walk and touch whatever objects are permitted. **Step 4:** The participant reads the perspicuous instruction that appears at the top of the screen. This instruction describes the task the user is expected to fulfill in the first level. Level 1 opening instruction: “Try to approach the spider” (Figure 4). **Step 5:** The participant executes the requested task. The aim of the first (easy) level is to allow users to observe the phobic object, to familiarize with it and thus exercise their response and composure when it appears in their personal space. In Level 1, the participant must: [a] Observe the virtual room: this helps users feel comfortable in the realm of the virtual environment, control their steps and learn how to synchronize their physical movements with those of their virtual body. [b] Approach the spider in their own time whilst keeping their composure: the participant can walk towards the table, where a static white spider lies; they can approach the spider whenever they feel ready and confident in themselves. The spider does not move at all, which gives users confidence and a strong sense of control. **Step 6:** The participant reads the final instruction that appears at the top of the screen. This instruction congratulates them on successfully completing the task and marks the end of the first level. Level 1 final message: “Congratulations!” (Figure 4). A demonstration video of the system is available here: [youtube.com/watch?v=Fcj9uE_wv0I](https://www.youtube.com/watch?v=Fcj9uE_wv0I).




Figure 3: (*top*) The first virtual environment in which the spider is on the table and the participant can approach it and touch it; (*bottom*) The second virtual environment in which the spider is appearing to hang and the participant can touch her and hit her.

Level 02: Step 1: The participant remains in the designated area without taking the VRG off, since Level 2 starts automatically once level 1 is completed. **Step 2:** The patient enters a new virtual room; each level has its own virtual room. **Step 3:** The patient reads the perspicuous instruction that appears at the top of the screen. This instruction describes the task the user is expected to complete during the second level. Level 2 opening instruction: “Try hitting the spider 3 times”. (Figure 5) **Step 4:** The participant executes the requested assignment. The aim of the second (hard) level is to allow users to touch the stimulus and learn how to remain calm, as well as confront the spider when it appears in their personal space. In Level 2 the participant must: [a] approach the spider in their own time whilst keeping their composure: in this level, a black spider is hanging from the ceiling using its web, which gives it the ability to swing whenever the user hits it; the participant can walk toward the spider when they feel ready and confident in themselves. The spider does not move unless the user hits it; this gives them time to relax and realize that the spider will not approach them unexpectedly; therefore, they have full control of the situation. [b] The participant is requested to hit the spider three times in the virtual world and manage to stay close; the width of the spider’s swing depends on the power of each hit, which increases the patient’s sense of control. They have to repeat the same action three times, so as to gradually realize that the spider cannot truly hurt them but instead, they can repel it if they want to; the patient can even move a step back whenever the spider swings towards them; yet, it is crucial that they keep their composure, get back to the hitting position and continue (Figure 5). **Step 5:** After




Figure 4. The participant has to execute an assignment: (*top*) The first aim is to try to approach the spider; (*bottom*) Once the participant has managed to get close the assignment will be completed.




Figure 5. The participant enters a different virtual room; (*top*) The participant first sees a hanging spider; Then a phrase appears which says “TRY HITTING THE SPIDER 3 TIMES”; (*bottom*) The participant hits the spider.

completing the task, the participant reads the final instruction that appears at the top of the screen. This instruction congratulates them on successfully completing the task and marks the end of the second level: “Congratulations!”. A demonstration video is available here: [youtube.com/watch?v=qDBbSoOrUKY](https://www.youtube.com/watch?v=qDBbSoOrUKY).

4 Results

As we can see in Table 1, the clinicians reportedly mentioned that action is crucial for treating Specific Phobias, since 80% answered “MAYBE YES” or “YES”. In Table 2, we can see that 80% of them answered “YES” or “MAYBE YES” to whether they think that the proposed system with MRC interaction can simulate real-life experiences. Last but not least, in Table 3, we can see a reduction of positive results, since 50% of the participating clinicians answered “MAYBE YES” or “YES” to whether coping mechanisms acquired in a virtual environment can be transferred to the real world. However, only 2 out of 20 answered “YES”.

Thereby, we conclude that not only is interaction necessary for the treatment process but also that the proposed system can simulate an appropriate interaction between the stimulus and the user through the VR simulation; however, according to the 20

clinicians, it is yet unclear whether virtual reality can actually assist patients in real life, despite the full body interaction.

Overall, the participants were initially thrilled with the opportunity to interact with the virtual environment of the room. What was particularly interesting is that participants completed their tasks easily in the first level, where they were not permitted to interact with the spider; nonetheless, during level 2, where participants were expected to interact with the stimulus, they were amazed by the possibility of interacting with the spider in such a straightforward way. Due to that, we can deduce that interaction with the feared object in the spectrum of a virtual environment could be far more efficient than the sole habituation of patients' anxiety through vivid virtual representations of the stimulus, i.e. images and videos. Also, even though participants were not familiar with the technology of the proposed system, no issues emerged in setting it up. As we mentioned above, the only prerequisite for the setup of the system is a 5 m² space for outlining the Action Area. Nevertheless, no issues were encountered during the rest of the process (Figure 6).




Table 1. The answers from the question Q1: “Do you believe that Exposure Therapy could benefit from the addition of interaction?”.




Table 2. The answers from the question Q2: “Do you think that interaction with the Motion Recognition Camera in the Virtual Environment can give real life experience?”.




Table 3. The answers from the question Q3: “Do you believe that coping mechanisms learnt in virtual situations and environments can be transferred to real-life situations?”.



Figure 6. Photos from the trial in the Hospital: (*top*) Here the participant tries to hit the spider with her punches; (*bottom*) Here the participant sees his hands in the virtual room.

5 Discussion

The present study investigated the possibility of expanding current ET treatments by adding another step, a new cycle of sessions in the procedure, the one we refer to as “Action Therapy”. The main intention of that step is to complement the visual stimuli already used in VRET by enhancing the confrontational capabilities of the patient towards their phobia during the treatment process. Our hypothesis is that the experience patients gain through the VR simulation could be more constructive and contribute significantly to their progress.

Thus, to determine whether the proposed Action Therapy could improve ET and be beneficial to patients, we tested the proposed system on a group of 20 people consisting of professionals in the field of psychiatry. The aim of this trial was to introduce them to the system and, based on their experience, get their feedback on whether or not it could be beneficial for patients and add to their training.

Our hypothesis has been confirmed to a large extent since the majority of clinicians found the addition of action to the existing treatment procedure promising for patients. They also partially confirmed that coping mechanisms acquired during VR could be utilized in real-life; however, half of the participants presume that VR has to further improve before becoming an apprentice in the treatment of mental illnesses.

The system we used included VRG, as well as a Motion Recognition Camera and, during the procedure, the participants had the chance to interact within a virtual environment specifically designed for arachnophobes. In spite of the promising feedback we received, there are limitations to take into consideration. Regarding the procedure, the trial was conducted on only a small group of 20 clinicians; therefore, the data was not extensive.

Another issue worthy of discussion is our choice to use the Motion Recognition Camera instead of other motion recognition tools, in order to add interaction to VR systems. Thanks to technological advances, there are various commercial motion tracking devices financially accessible (from the Motion Recognition Camera to Wearable Body Tracking Sensors, like gloves and suits with gyroscope trackers or accelerometer trackers, or Hand Controllers). The main reason we did not choose to use Wearable Motion Tracking Sensors is because they limit the interaction capability of the patient’s body within the virtual environment, thus potentially rendering their movements unnatural, i.e. they have to press certain buttons instead of using their hands, so as to touch objects (Chen, Qiu, & Ahn, 2017; Da Gama, Chaves, Fallavollita, Figueiredo, & Teichrieb, 2019; Wang & Yan, 2017). Nevertheless, the MRC has some limitations as well, since it cannot track movements in all body parts (e.g. the fingers) due to its range (0.6-8.0m). However, the purpose of this study was not to extensively examine the differences between different types of motion tracking sensors, but to determine whether interaction, in general, could improve existing treatment protocols.

Finally, another interesting aspect of our study was that MRCs offer a remarkable feature: the tracking of the patient’s physical movements is not only useful for placing their body in the virtual environment, but also for dispatching the information obtained by their movements in real time. Measurements of the patient’s movements as they executed the exercises can help draw further conclusions about the patient’s performance, i.e. whether the patient completed the task effortlessly or not. They could

also diagnose patients using the same system by comparing their data to those of healthy individuals.

6 Conclusion

Up to now, there has been no experimental indication stemming from clinical trial results that VRET is more effective than in VIVO ET. Nevertheless, previous studies have found them equally realistic for patients. Considering the technical issues of In VIVO ET such as affordability, the time needed to implement it, as well as the fact that it is impossible for clinicians to acquire every phobic stimulus, VRET seems to be at least as helpful as In VIVO ET (Maples-Keller, Bunnell, Kim, & Rothbaum, 2017). Acquisition costs of VR systems have dropped significantly, making it possible for VRET to be applied in a larger scale in clinicians' offices. Following, we propose that biomedical laboratories and tech-oriented research centers focus more on technologies that fit in the needs of anxiety disorder field.

Overall, VR applications related to anxiety disorders are still on an early stage of development, without any solid foundations built yet. Currently, this field seeks to adapt itself to the newly introduced technological tools (D. Freeman et al., 2017; Morina, Ijntema, Meyerbröker, & Emmelkamp, 2015; Suh & Prophet, 2018; Turner & Casey, 2014); at the same time, the technological tools themselves, also need to adapt to the respective safety regulations (Blakey & Abramowitz, 2016). Our aim, through our research, focuses on encouraging the creation of technological tools precisely for the needs of this industry.

Finally, we suggest the proposed system could be an effective tool in other areas. By modifying the scenarios, it could be useful for people with development disorders to control their movements and improve their social skills or in the area of forensic psychiatry, i.e. panic attacks, heart disease, epilepsy, or to people who take drugs with large psychological effects. The instruments currently in use are mainly subjective questionnaires. By adding cardiac rhythm and sweating sensors to our platform we can offer an objective evaluation based primarily on the immediate bodily responses towards the stimuli without the interference of the possible manipulation of the questionnaires by the persons assessed.

Acknowledgments

We would like to thank the executive managers and members of the staff of the General and Oncology Hospital of Kifissia for agreeing to provide us with the opportunity to present our work to them.

At this point, we would like to discuss the expertise of the healthcare facility members we used as evaluators of our system a bit further. The medical staff of the facility we chose consists of experienced, well-trained, highly educated psychiatrists and healthcare attendants, who agreed to test our proposed system and provide their substantial feedback on its efficiency, as well as methods of enhancing it in the future. All of the members have graduated from reputable medical universities and have frequently assisted patients in treating their obsessive/irrational thought and behavioral patterns through exposure therapy sessions. Another important aspect of our preference in this

particular psychiatric department is their constant quest for innovative diagnostic and therapeutic techniques. The VR system we propose is an original method; subsequently, they were highly interested in the new methods in which exposure therapy can be enhanced through VR technology. Taking the aforementioned facts into consideration, it is clear that our choice in this particular psychiatric department was everything but coincidental; we strongly believe that the clinic's specialists were the most fitting to evaluate the efficacy of our system in view of their experience and education.

References

- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders (5th ed.)*. *American Journal of Psychiatry*. <https://doi.org/10.1176/appi.books.9780890425596.744053>
- American Psychiatric Association. (2013). *DSM-V. Manual Diagnóstico E Estatístico De Transtornos Mentais - DSM-V*. <https://doi.org/10.1176/appi.books.9780890425596.744053>
- Bandelow, B., & Michaelis, S. (2015). Epidemiology of anxiety disorders in the 21st century. *Dialogues in Clinical Neuroscience*, 17(3), 327–335. <https://doi.org/10.1016/j.siny.2015.10.004>
- Blakey, S. M., & Abramowitz, J. S. (2016). The effects of safety behaviors during exposure therapy for anxiety: Critical analysis from an inhibitory learning perspective. *Clinical Psychology Review*. <https://doi.org/10.1016/j.cpr.2016.07.002>
- Bun, P., Gorski, F., Grajewski, D., Wichniarek, R., & Zawadzki, P. (2016). Low - Cost Devices Used in Virtual Reality Exposure Therapy. In *Procedia Computer Science*. <https://doi.org/10.1016/j.procs.2017.01.158>
- Cardoş, R. A. I., David, O. A., & David, D. O. (2017). Virtual reality exposure therapy in flight anxiety: A quantitative meta-analysis. *Computers in Human Behavior*. <https://doi.org/10.1016/j.chb.2017.03.007>
- Carlin, A. S., Hoffman, H. G., & Weghorst, S. (1997). Virtual reality and tactile augmentation in the treatment of spider phobia: A case report. *Behaviour Research and Therapy*. [https://doi.org/10.1016/S0005-7967\(96\)00085-X](https://doi.org/10.1016/S0005-7967(96)00085-X)
- Chen, J., Qiu, J., & Ahn, C. (2017). Construction worker's awkward posture recognition through supervised motion tensor decomposition. *Automation in Construction*. <https://doi.org/10.1016/j.autcon.2017.01.020>
- Da Gama, A. E. F., Chaves, T. de M., Fallavollita, P., Figueiredo, L. S., & Teichrieb, V. (2019). Rehabilitation motion recognition based on the international biomechanical standards. *Expert Systems with Applications*. <https://doi.org/10.1016/j.eswa.2018.09.026>
- Emmelkamp, P. M. G., Krijn, M., Hulsbosch, A. M., de Vries, S., Schuemie, M. J., & van der Mast, C. A. P. G. (2002). Virtual reality treatment versus exposure in vivo: a comparative evaluation in acrophobia. *Behaviour Research and Therapy*. [https://doi.org/10.1016/S0005-7967\(01\)00023-7](https://doi.org/10.1016/S0005-7967(01)00023-7)
- Freeman, D., Haselton, P., Freeman, J., Spanlang, B., Kishore, S., Albery, E., ... Nickless, A. (2018). Automated psychological therapy using immersive virtual reality for treatment of fear of heights: a single-blind, parallel-group, randomised

- controlled trial. *The Lancet Psychiatry*. [https://doi.org/10.1016/S2215-0366\(18\)30226-8](https://doi.org/10.1016/S2215-0366(18)30226-8)
- Freeman, D., Reeve, S., Robinson, A., Ehlers, A., Clark, D., Spanlang, B., & Slater, M. (2017). Virtual reality in the assessment, understanding, and treatment of mental health disorders. *Psychological Medicine*. <https://doi.org/10.1017/S003329171700040X>
- Kerns, C. M., Read, K. L., Klugman, J., & Kendall, P. C. (2013). Cognitive behavioral therapy for youth with social anxiety: Differential short and long-term treatment outcomes. *Journal of Anxiety Disorders*, 27(2), 210–215. <https://doi.org/10.1016/j.janxdis.2013.01.009>
- Kessler, R. C., Berglund, P., Demler, O., Jin, R., Merikangas, K. R., & Walters, E. E. (2005). Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the national comorbidity survey replication. *Archives of General Psychiatry*. <https://doi.org/10.1001/archpsyc.62.6.593>
- Krebs, G., de la Cruz, L. F., Monzani, B., Bowyer, L., Anson, M., Cadman, J., ... Mataix-Cols, D. (2017). Long-Term Outcomes of Cognitive-Behavioral Therapy for Adolescent Body Dysmorphic Disorder. *Behavior Therapy*, 48(4), 462–473. <https://doi.org/10.1016/j.beth.2017.01.001>
- Krijn, M., Emmelkamp, P. M. G., Biemond, R., De Wilde De Ligny, C., Schuemie, M. J., & Van Der Mast, C. A. P. G. (2004). Treatment of acrophobia in virtual reality: The role of immersion and presence. *Behaviour Research and Therapy*. [https://doi.org/10.1016/S0005-7967\(03\)00139-6](https://doi.org/10.1016/S0005-7967(03)00139-6)
- Maples-Keller, J. L., Bunnell, B. E., Kim, S. J., & Rothbaum, B. O. (2017). The use of virtual reality technology in the treatment of anxiety and other psychiatric disorders. *Harvard Review of Psychiatry*. <https://doi.org/10.1097/HRP.0000000000000138>
- Mayo-Wilson, E., & Montgomery, P. (2013). Media-delivered cognitive behavioural therapy and behavioural therapy (self-help) for anxiety disorders in adults. *Cochrane Database of Systematic Reviews*. <https://doi.org/10.1002/14651858.CD005330.pub4>
- Morina, N., Brinkman, W.-P., Hartanto, D., & Emmelkamp, P. M. G. (2014). Sense of presence and anxiety during virtual social interactions between a human and virtual humans. *PeerJ*. <https://doi.org/10.7717/peerj.337>
- Morina, N., Ijntema, H., Meyerbröker, K., & Emmelkamp, P. M. G. (2015). Can virtual reality exposure therapy gains be generalized to real-life? A meta-analysis of studies applying behavioral assessments. *Behaviour Research and Therapy*. <https://doi.org/10.1016/j.brat.2015.08.010>
- National Institute of Mental Health - Anxiety Disorder. (n.d.). Retrieved from <https://www.nimh.nih.gov/health/statistics/any-anxiety-disorder.shtml>
- National Institute of Mental Health - Mental Illness. (n.d.).
- Ollendick, T. H., Öst, L. G., Ryan, S. M., Capriola, N. N., & Reuterskiöld, L. (2017). Harm beliefs and coping expectancies in youth with specific phobias. *Behaviour Research and Therapy*, 91, 51–57. <https://doi.org/10.1016/j.brat.2017.01.007>
- Öst, L. G. (1989). One-session treatment for specific phobias. *Behaviour Research and Therapy*. [https://doi.org/10.1016/0005-7967\(89\)90113-7](https://doi.org/10.1016/0005-7967(89)90113-7)

- Otte, C. (2011). Cognitive behavioral therapy in anxiety disorders: Current state of the evidence. *Dialogues in Clinical Neuroscience*, *13*(4), 413–421. <https://doi.org/10.1186/1471-244X-14-S1-S1>
- Peñate, W., Fumero, A., Viña, C., Herrero, M., Marrero, R. J., & Rivero, F. (2017). A meta-analytic review of neuroimaging studies of specific phobia to small animals. *European Journal of Psychiatry*. <https://doi.org/10.1016/j.ejpsy.2016.12.003>
- Price, M., & Anderson, P. (2007). The role of presence in virtual reality exposure therapy. *Journal of Anxiety Disorders*. <https://doi.org/10.1016/j.janxdis.2006.11.002>
- Riddle-Walker, L., Veale, D., Chapman, C., Ogle, F., Rosko, D., Najmi, S., ... Hicks, T. (2016). Cognitive behaviour therapy for specific phobia of vomiting (Emetophobia): A pilot randomized controlled trial. *Journal of Anxiety Disorders*, *43*, 14–22. <https://doi.org/10.1016/j.janxdis.2016.07.005>
- Rizzo, A., Hartholt, A., Grimani, M., Leeds, A., & Liewer, M. (2014). Virtual reality exposure therapy for combat-related posttraumatic stress disorder. *Computer*. <https://doi.org/10.1109/MC.2014.199>
- Rothbaum, B. O., Rizzo, A. S., McDaniel, D. D., & Zanov, M. V. (2015). Virtual Reality Exposure Therapy. In *Encyclopedia of Mental Health: Second Edition*. <https://doi.org/10.1016/B978-0-12-397045-9.00196-8>
- Rozental, A., Shafraan, R., Wade, T., Egan, S., Nordgren, L. B., Carlbring, P., ... Andersson, G. (2017). A randomized controlled trial of Internet-Based Cognitive Behavior Therapy for perfectionism including an investigation of outcome predictors. *Behaviour Research and Therapy*, *95*, 79–86. <https://doi.org/10.1016/j.brat.2017.05.015>
- Rudaz, M., Ledermann, T., Margraf, J., Becker, E. S., & Craske, M. G. (2017). The moderating role of avoidance behavior on anxiety over time: Is there a difference between social anxiety disorder and specific phobia? *PLoS ONE*. <https://doi.org/10.1371/journal.pone.0180298>
- Scott, J., & Beck, A. T. (2008). Cognitive behavioural therapy. In *Essential Psychiatry, Fourth Edition*. <https://doi.org/10.1017/CBO9780511544125.029>
- Shiban, Y., Schelhorn, I., Pauli, P., & Mühlberger, A. (2015). Effect of combined multiple contexts and multiple stimuli exposure in spider phobia: A randomized clinical trial in virtual reality. *Behaviour Research and Therapy*, *71*, 45–53. <https://doi.org/10.1016/j.brat.2015.05.014>
- Society of Clinical Psychology. (2013). What is Exposure Therapy?
- Stinson, F. S., Dawson, D. a, Patricia Chou, S., Smith, S., Goldstein, R. B., June Ruan, W., & Grant, B. F. (2007). The epidemiology of DSM-IV specific phobia in the USA: results from the National Epidemiologic Survey on Alcohol and Related Conditions. *Psychological Medicine*, *37*(March), 1047–1059. <https://doi.org/10.1017/S0033291707000086>
- Suh, A., & Prophet, J. (2018). The state of immersive technology research: A literature analysis. *Computers in Human Behavior*. <https://doi.org/10.1016/j.chb.2018.04.019>
- Treanor, M., & Craske, M. G. (2015). Exposure Therapy. In *Encyclopedia of Mental Health: Second Edition*. <https://doi.org/10.1016/B978-0-12-397045-9.00024-0>

- Treatment - (ADAA) Anxiety and Depression Association of America. (n.d.). Retrieved from <https://www.adaa.org/finding-help/treatment>
- Turner, W. A., & Casey, L. M. (2014). Outcomes associated with virtual reality in psychological interventions: where are we now? *Clinical Psychology Review*. <https://doi.org/10.1007/s12517-015-2040-z>
- Twomey, C., O'Reilly, G., & Meyer, B. (2017). Effectiveness of an individually-tailored computerised CBT programme (Deprexis) for depression: A meta-analysis. *Psychiatry Research*. <https://doi.org/10.1016/j.psychres.2017.06.081>
- Wang, X., & Yan, K. (2017). Immersive human-computer interactive virtual environment using large-scale display system. *Future Generation Computer Systems*. <https://doi.org/10.1016/j.future.2017.07.058>
- Warwick, H., Reardon, T., Cooper, P., Murayama, K., Reynolds, S., Wilson, C., & Creswell, C. (2017). Complete recovery from anxiety disorders following Cognitive Behavior Therapy in children and adolescents: A meta-analysis. *Clinical Psychology Review*. <https://doi.org/10.1016/j.cpr.2016.12.002>
- Waters, A. M., Farrell, L. J., Zimmer-Gembeck, M. J., Milliner, E., Tiralongo, E., Donovan, C. L., ... Ollendick, T. H. (2014). Augmenting one-session treatment of children's specific phobias with attention training to positive stimuli. *Behaviour Research and Therapy*, 62, 107–119. <https://doi.org/10.1016/j.brat.2014.07.020>
- Wittchen, H.-U., Mühlig, S., & Beesdo, K. (2003). Mental disorders in primary care. *Dialogues in Clinical Neuroscience*, 5(2), 115–128. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/22034245> <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC3181625>
- Yiasemidou, M., de Siqueira, J., Tomlinson, J., Glassman, D., Stock, S., & Gough, M. (2017). “Take-home” box trainers are an effective alternative to virtual reality simulators. *Journal of Surgical Research*. <https://doi.org/10.1016/j.jss.2017.02.038>
- Society of Clinical Psychology. (2013). What is Exposure Therapy? *American Psychological Association: Division 12*, 12. Retrieved from <http://www.apa.org/ptsd-guideline/patients-and-families/exposure-therapy.aspx>