Benzodiazepine Withdrawal Symptoms

COMMON WITHDRAWAL SYMPTOMS

Abdominal pains and cramps

Agoraphobia Anxiety

Breathing difficulties Blurred vision

Changes in perception (faces distorting and inanimate

objects or surfaces moving)

Depression

Distended abdomen

Dizziness

Extreme lethargy
Fears (uncharacteristic)
Feelings of unreality
Flu-like symptoms
Heavy limbs
Heart palpitations

Hypersensitivity to light

Indigestion Insomnia Irritability

Lack of concentration Lack of coordination Loss of balance Loss of memory

Muscular aches and pains

Nausea Nightmares Panic attacks

Rapid mood changes (crying one moment, laughing the

next)

Restlessness Severe headaches

Shaking Seeing spots Sore eyes Sweating

Tightness in the chest

Tightness in the head (feeling of a band around the head)

LESS COMMON WITHDRAWAL SYMPTOMS

Aching jaw

Craving for sweet food

Constipation

Depersonalisation (a feeling of not knowing who

you are) Diarrhoea

Difficulty swallowing

Feelings of the ground moving Hallucinations (auditory and visual)

Hyperactivity

Hypersensitivity to sound

Incontinence or frequency and urgency (needing to

urinate often and in a hurry)

Increased saliva

Loss or change in appetite

Loss of taste, or changes in taste (e.g. a metallic taste

in the mouth)

Menstrual problems (painful periods, irregular

periods or cessation of periods)

Morbid thoughts

Numbness in any part of the body Outbursts of rage and aggression

Paranoia Painful scalp

Persistent, unpleasant memories

Pins and needles

Rapid changes in body temperature Sexual problems (changes in libido)

Skin problems (dryness, itchiness, rashes, slow healing hoils)

ing, boils)

Sore mouth and tongue (ulcers, cracked lips, cold

sores)

Speech difficulties Suicidal thoughts

Tinnitus (buzzing or ringing in the ears)

Unusually emotionally sensitive (e.g. unable to watch

the news on television) Vaginal discharge

Vomiting

VOITHCHIE

Weight loss or gain

RARE WITHDRAWAL SYMPTOMS

Blackouts (this is rare with low dose use, but less rare when large doses have been taken)

Bleeding from the nose

Burning sensation around the mouth

Craving for pills

Discharge from the breasts

Burning along the spine

Falling hair

Haemorrhoids

Hypersensitivity to touch

Menorrhagia (bleeding between menstrual cycles)

Rectal bleeding Sinus pain Seizures (fits)

Sensitive or painful teeth

Swollen breasts

