


Total Synthesis of Tagetitoxin


He, C.; Chu, H.; Stratton, T. P.; Kossler, D.; Eberle, K. J.; Flood, D. T.; Baran, P. S.
J. Am. Chem. Soc. **2020**, *142*, 13683–13688.




1–7



8–10



1




2

- 1) **1**, amberlyst 15, Δ
- 2) **2**, Et₃N *then* 2,2,2-trifluoroethanol, TBAF
- 3) methylene blue, O₂, h ν *then* Me₂S, SiO₂
- 4) 4-cyanobenzoyl chloride, DMAP
- 5) CeCl₃·7H₂O, NaBH₄
- 6) 1,1'-thiocarbonyldiimidazole
- 7) BHT, 115 °C *then* *p*-TsOH, H₂O


3) Please provide a mechanism.

5) Please name the reaction.

7) Please classify the reaction.




3



4

- 8) **3**, Et₃N
- 9) OsO₄, NMO, citric acid *then* HCl
- 10) **4**, AcOH, Δ



9) Hint: The most labile protective group is lost.



(+)-tagetitoxin



11–15



- 11) TfOH, anisole
- 12) *n*-Bu₃SnH, AIBN *then* acetone, *p*-TsOH
- 13) MeOH, Et₃N *then* **5**, DBN; separation of diastereomers
- 14) SeO₂
- 15) TMSOK, H₂O *then* MeONH₂·HCl

11) Hint: Monodeprotection.

13) Hint: 2 eq of MeOH are consumed.