

Write your name here

Surname

Other names

**Pearson Edexcel**  
**Level 3 GCE**

Centre Number

--	--	--	--	--

Candidate Number

--	--	--	--	--

# Further Mathematics

**Advanced**  
**Further Mathematics Option 1**  
**Paper 3: Further Mechanics 1**  
**Further Mathematics Option 2**  
**Paper 4: Further Mechanics 1**

Sample Assessment Material for first teaching September 2017

**Time: 1 hour 30 minutes**

Paper Reference

**9FM0/3C**  
**9FM0/4C**

**You must have:**

Mathematical Formulae and Statistical Tables, calculator

Total Marks

--

**Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for algebraic manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.**

## Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided  
– *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Answers should be given to three significant figures unless otherwise stated.

## Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 8 questions in this question paper. The total mark for this paper is 75.
- The marks for **each** question are shown in brackets  
– *use this as a guide as to how much time to spend on each question.*

## Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

S54442A

©2017 Pearson Education Ltd.

1/1/1/1/



Pearson



**Question 1 continued**

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 1 is 6 marks)

2. A parcel of mass 5 kg is projected with speed  $8 \text{ m s}^{-1}$  up a line of greatest slope of a fixed rough inclined ramp.

The ramp is inclined at angle  $\alpha$  to the horizontal, where  $\sin \alpha = \frac{1}{7}$

The parcel is projected from the point  $A$  on the ramp and comes to instantaneous rest at the point  $B$  on the ramp, where  $AB = 14 \text{ m}$ .

The coefficient of friction between the parcel and the ramp is  $\mu$ .

In a model of the parcel’s motion, the parcel is treated as a particle.

(a) Use the work-energy principle to find the value of  $\mu$ . (5)

(b) Suggest one way in which the model could be refined to make it more realistic. (1)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

**Question 2 continued**

DO NOT WRITE IN THIS AREA

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

**(Total for Question 2 is 6 marks)**

3. A particle of mass  $m$  kg lies on a smooth horizontal surface.

Initially the particle is at rest at a point  $O$  between two fixed parallel vertical walls.

The point  $O$  is equidistant from the two walls and the walls are 4 m apart.

At time  $t = 0$  the particle is projected from  $O$  with speed  $u$   $\text{ms}^{-1}$  in a direction perpendicular to the walls.

The coefficient of restitution between the particle and each wall is  $\frac{3}{4}$

The magnitude of the impulse on the particle due to the first impact with a wall is  $\lambda mu$  N s.

(a) Find the value of  $\lambda$ . (3)

The particle returns to  $O$ , having bounced off each wall once, at time  $t = 7$  seconds.

(b) Find the value of  $u$ . (5)



4.




Figure 1

Figure 1 represents the plan view of part of a horizontal floor, where  $AB$  and  $BC$  are perpendicular vertical walls.

The floor and the walls are modelled as smooth.

A ball is projected along the floor towards  $AB$  with speed  $u \text{ m s}^{-1}$  on a path at an angle of  $60^\circ$  to  $AB$ . The ball hits  $AB$  and then hits  $BC$ .

The ball is modelled as a particle.

The coefficient of restitution between the ball and wall  $AB$  is  $\frac{1}{\sqrt{3}}$

The coefficient of restitution between the ball and wall  $BC$  is  $\sqrt{\frac{2}{5}}$

(a) Show that, using this model, the final kinetic energy of the ball is 35% of the initial kinetic energy of the ball.

(8)

(b) In reality the floor and the walls may not be smooth. What effect will the model have had on the calculation of the percentage of kinetic energy remaining?

(1)



Question 4 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Lined writing area for the question response, consisting of multiple horizontal lines.

Question 4 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

**Question 4 continued**

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

**(Total for Question 4 is 9 marks)**

5. A car of mass 600 kg is moving along a straight horizontal road.

At the instant when the speed of the car is  $v \text{ m s}^{-1}$ , the resistance to the motion of the car is modelled as a force of magnitude  $(200 + 2v) \text{ N}$ .

The engine of the car is working at a constant rate of 12 kW.

- (a) Find the acceleration of the car at the instant when  $v = 20$

(4)

Later on the car is moving up a straight road inclined at an angle  $\theta$  to the horizontal,

where  $\sin \theta = \frac{1}{14}$

At the instant when the speed of the car is  $v \text{ m s}^{-1}$ , the resistance to the motion of the car from non-gravitational forces is modelled as a force of magnitude  $(200 + 2v) \text{ N}$ .

The engine is again working at a constant rate of 12 kW.

At the instant when the car has speed  $w \text{ m s}^{-1}$ , the car is decelerating at  $0.05 \text{ m s}^{-2}$ .

- (b) Find the value of  $w$ .

(5)





**Question 6 continued**

Lined writing area for the answer.

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA





**Question 6 continued**

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

**(Total for Question 6 is 9 marks)**

7. A particle  $P$  of mass  $m$  is attached to one end of a light elastic string of natural length  $a$  and modulus of elasticity  $3mg$ .

The other end of the string is attached to a fixed point  $O$  on a ceiling.

The particle hangs freely in equilibrium at a distance  $d$  vertically below  $O$ .

- (a) Show that  $d = \frac{4}{3}a$ . (3)

The point  $A$  is vertically below  $O$  such that  $OA = 2a$ .

The particle is held at rest at  $A$ , then released and first comes to instantaneous rest at the point  $B$ .

- (b) Find, in terms of  $g$ , the acceleration of  $P$  immediately after it is released from rest. (3)

- (c) Find, in terms of  $g$  and  $a$ , the maximum speed attained by  $P$  as it moves from  $A$  to  $B$ . (5)

- (d) Find, in terms of  $a$ , the distance  $OB$ . (3)

**Question 7 continued**

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Lined area for student response.

**Question 7 continued**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

**Question 7 continued**

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

**(Total for Question 7 is 14 marks)**

8. A particle  $P$  of mass  $2m$  and a particle  $Q$  of mass  $5m$  are moving along the same straight line on a smooth horizontal plane.

They are moving in opposite directions towards each other and collide directly.

Immediately before the collision the speed of  $P$  is  $2u$  and the speed of  $Q$  is  $u$ .

The direction of motion of  $Q$  is reversed by the collision.

The coefficient of restitution between  $P$  and  $Q$  is  $e$ .

- (a) Find the range of possible values of  $e$ . (8)

Given that  $e = \frac{1}{3}$

- (b) show that the kinetic energy lost in the collision is  $\frac{40mu^2}{7}$ . (5)

- (c) Without doing any further calculation, state how the amount of kinetic energy lost in the collision would change if  $e > \frac{1}{3}$  (1)







**Question 8 continued**

Lined area for writing answers.

**(Total for Question 8 is 14 marks)**

**TOTAL FOR PAPER IS 75 MARKS**

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA