

Beyond antiplatelets: The role of glycoprotein VI in ischemic stroke

Isuru Induruwa¹, Stephanie M Jung² and Elizabeth A Warburton¹

International Journal of Stroke

0(0) 1–8

© 2016 World Stroke Organization

Reprints and permissions:

sagepub.co.uk/journalsPermissions.nav

DOI: 10.1177/1747493016654532

wso.sagepub.com



Abstract

Background: Platelets are essential to physiological hemostasis or pathological thrombus formation. Current antiplatelet agents inhibit platelet aggregation but leave patients at risk of systemic side-effects such as hemorrhage. Newer therapeutic strategies could involve targeting this cascade earlier during platelet adhesion or activation via inhibitory effects on specific glycoproteins, the thrombogenic collagen receptors found on the platelet surface.

Aims: Glycoprotein VI (GPVI) is increasingly being recognized as the main platelet-collagen receptor involved in arterial thrombosis. This review summarizes the crucial role GPVI plays in ischemic stroke as well as the current strategies used to attempt to inhibit its activity.

Summary of review: In this review, we discuss the normal hemostatic process, and the role GPVI plays at sites of atherosclerotic plaque rupture. We discuss how the unique structure of GPVI allows for its interaction with collagen and creates downstream signaling that leads to thrombus formation. We summarize the current strategies used to inhibit GPVI activity and how this could translate to a clinically viable entity that may compete with current antiplatelet therapy.

Conclusion: From animal models, it is clear that GPVI inhibition leads to an abolished platelet response to collagen and reduced platelet aggregation, culminating in smaller arterial thrombi. There is now an increasing body of evidence that these findings can be translated into the development of a bleeding free pharmacological entity specific to sites of plaque rupture in humans.

Keywords

Stroke, platelets, glycoprotein VI, antiplatelets

Received: 23 February 2016; accepted: 3 May 2016

Introduction

Ischemic stroke is a worldwide leading cause of disability and death.¹ An important and distinct subtype of ischemic stroke is due to the acute rupture of atherosclerotic plaques seen in large artery disease.² The mechanism of cerebral hypoperfusion involves platelets binding to exposed sub-endothelial collagen (adhesion), resulting in platelet activation, aggregation, and thrombus formation. Unstable thrombi can detach and travel to cerebral vessels causing stroke. As a result, the current management of acute stroke involves using antiplatelet agents such as aspirin and clopidogrel that inhibit platelet activation/aggregation, but often at the risk of off-target adverse effects such as hemorrhage.^{3,4}

Novel therapeutic strategies could involve targeting this cascade earlier during platelet adhesion or activation via inhibitory effects on specific platelet glycoproteins, the thrombogenic collagen receptors on their surfaces. Glycoprotein VI (GPVI) is one such

crucial transmembrane collagen receptor and pharmacological inhibition of GPVI, in order to stop pathological thrombus formation specific to the site of vessel injury, is currently being explored.

Hemostasis – clotting and thrombus formation

Platelets are anucleate cells derived from megakaryocytes. They contain unique cytoplasmic structures, α - and dense granules that can rapidly release their contents upon activation, promoting thrombus formation.

¹Department of Clinical Neurosciences, Box 83, Cambridge University Biomedical Campus, Cambridge, UK


²Department of Biochemistry, University of Cambridge, Cambridge, UK

Corresponding author:

Isuru Induruwa, Department of Clinical Neurosciences, Cambridge University Hospitals NHS Foundation Trust, R3 Addenbrookes Hospital, Cambridge, CB2 0QQ, UK.

Email: ipinduruwa@googlemail.com

Figure 1. Collagen-binding receptors (GPIb, GPVI, and integrin $\alpha 2\beta 1$) involved in platelet adhesion and activation. Sub-endothelial collagen exposed upon vessel injury binds to von Willebrand factor (vWF) in the blood (step 1). Platelets become tethered to collagen fibers by their vWF receptor GPIb—a weak interaction—so platelets transiently bind and detach, moving along the collagen (step 2). Platelets become firmly attached when their collagen receptors GPVI dimer and integrin $\alpha 2\beta 1$ binds to collagen; signaling through either GPVI or GPIb converts integrin $\alpha 2\beta 1$ to its high affinity form (step 3). GPVI engagement with collagen initiates a signaling cascade that culminates in platelet activation, spreading, and granule contents release (step 4), recruiting other platelets and forming a thrombus (steps 5 and 6). Integrin $\alpha IIb\beta 3$ (not shown) becomes activated through inside-out-signaling, enabling it to bind fibrinogen, through which inter-platelet bridges can be formed, allowing thrombus propagation.



Under normal hemostasis numerous protective barriers to thrombus formation exist to contain it to the injured site. This includes the continuous lining of endothelium that prevents platelets coming into contact with the prothrombotic sub-endothelial matrix,⁵ expression of ectonucleoside triphosphate diphosphohydrolase (CD39/ENTPD1)⁶ and secretion of prostacyclin (PGI₂) and nitric oxide.

There are four main platelet glycoprotein receptors that participate in the platelet–collagen interaction and facilitate thrombus formation. The GPIb-IX-V complex (GPIb α , GPIb β , GPIX and GPV) binds to von Willebrand factor (VWF) immobilized on collagen. Both GP Ia/IIa (integrin $\alpha 2\beta 1$) and GPVI bind directly to exposed collagen. GPIIb/IIIa (integrin $\alpha IIb\beta 3$) are converted to their high-affinity forms via inside-out signaling in activated platelets, enabling them to bind free fibrinogen and VWF.


Fibrous collagen is the ligand for both GPVI and integrin $\alpha 2\beta 1$. Each collagen monomer comprises three 1000 amino acid polypeptides arranged in a triple helix, and bundles of these monomers align to form collagen fibers.⁷ With atherosclerotic plaque rupture, the sub-endothelial fibrillar collagens (types I and III) are exposed to the blood stream. Under high shear

(arterial flow), VWF becomes immobilized on the exposed collagen fibers and binds with the GPIb-IX-V complex^{8–10} (Figure 1). Platelets translocate on the collagen surface until firmly arrested through almost simultaneous binding with integrin $\alpha 2\beta 1$ and GPVI. GPVI engagement with collagen (via GPVI dimers) initiates a signaling cascade leading to platelet activation and inside-out signaling (and intracellular calcium mobilization) that converts low affinity forms of integrins $\alpha IIb\beta 3$ and $\alpha 2\beta 1$ into their active forms, as well as further clustering of GPVI receptors.⁸

Activated platelets rapidly synthesize thromboxane A₂ (TxA₂) and secrete this, along with the contents of their alpha (fibrinogen, P-selectin, and VWF multimers) and dense granules (ADP).⁹ An increase in intracellular calcium in response to ADP and TxA₂ induces platelet shape change to an irregular shape with multiple filipoidal surfaces. This facilitates the formation of a close structure of folded platelets in the platelet plug.

Now non-activated platelets are recruited into the growing thrombus, also becoming activated by ADP and TxA₂. GPVI-activated platelets provide a pro-coagulant surface for the generation of thrombin—separately through the coagulation cascade by

Figure 2. Structures of GPVI monomer and dimer. The extracellular domain of each monomer comprises two IgG domains (D1 and D2) and a mucin-like Ser/Thr-rich domain connecting D1/D2 to the transmembrane domain. The binding site recognizing the GPO triplets of collagen, resides in D1. Each monomer is non-covalently associated through a salt bridge with FcR γ . FcR γ is a disulphide-linked dimer, each chain containing an ITAM sequence, which when phosphorylated binds to the tyrosine kinase Syk. The phosphorylated Syk now initiates signaling. The short intracellular domain of GPVI contains a basic domain that binds to calmodulin; a proline-rich domain that binds to Src kinases Fyn and Lyn, which participate in phosphorylation of ITAM; and a C-terminal tail.



releasing several substances like FV, FXIII, fibrinogen, and Protein S. This further activates platelets and converts fibrinogen to fibrin, resulting in a strong mesh structure.

Glycoprotein VI and its structure

Glycoprotein VI, a 62-kDA transmembrane glycoprotein exclusively expressed in platelets and megakaryocytes, is associated with an immunoreceptor tyrosine-based-activation motif (ITAM) containing signaling subunit FcR γ (Figure 2).¹⁰ Its gene is mapped to 19q13.4 of the human genome. GPVI contains two IgG-like extracellular domains (D1 and D2) linked by a peptide strand; D2 is connected to the transmembrane domain via a glycosylated stem; and its 51 amino acid cytoplasmic tail is required for signal transmission.^{11,12}

GPVI exists in both monomeric and back-to-back dimeric forms on the platelet surface¹³ and forms a non-covalently linked complex with the Fc receptor γ -chain (FcR γ) via a salt bridge formed between their transmembrane domains. FcR γ itself is also a dimer,

held together by covalent links and is essential for GPVI expression, as FcR γ knockout mice do not express GPVI.¹⁴ Dimerization or multimerization of GPVI was first suggested by Berlanga et al.¹⁵ but proof of the actual existence on platelets was first provided by Jung et al.¹⁶ Later, they found that constitutive dimers of GPVI make up of about 20% of the total GPVI^{13,17} in resting platelets of normal individuals and platelet activation increases the number of dimers.¹⁷ Only GPVI dimer has high affinity for collagen, while the monomer binds weakly if at all. Inhibition by dimer specific antibody m-Fab-F markedly inhibited collagen-induced platelet aggregation, highlighting its role as the functional form of GPVI.

Interaction with collagen and signaling

So how does GPVI interact with the fibrous subendothelial collagen exposed upon vessel injury to induce platelet activation? Much of this data has been gained by using snake venom-derived convulxin and CRP (collagen-related peptide) in vitro, which activate platelets in a similar way to collagen.^{7,18} CRP, a triple

helical peptide containing 10 glycine–proline–hydroxyproline (GPO) sequences, is an especially potent platelet agonist specific for GPVI. GPVI was suggested to bind to the GPO sequences in collagen. Smethurst et al.¹⁹ later found that the minimum recognition motif for GPVI is one GPO.

Using the crystal structure of GPVI and a docking model with CRP, Horii et al.²⁰ found that GPVI formed back-to-back dimers, each containing a shallow groove on the D1 domain that is perpendicular to the collagen triple helix, which fits precisely into this 5.5-nm gap.

Loyau et al.²¹ proposed that at least two GPVI dimer-FcR γ complexes are required to create a working signaling unit, as the FcR γ chain is also essential for the function of GPVI and each chain contains an ITAM. Upon GPVI–collagen binding, the Tyr residues of the ITAM sequence become phosphorylated, initiating signal transduction. PCL γ 2 is subsequently activated, leading to eventual activation of protein kinase C, resulting in calcium mobilization, degranulation and GPIIb/IIIa activation and the start of the next step in the process—platelet aggregation.

GPVI shedding and receptor downregulation

An important consequence of GPVI downstream signaling, which serves to limit thrombus growth, is antibody or metalloproteinase (MMP) induced shedding of a 55 kDa GPVI ectodomain into the blood stream, leaving a 10 kDa remnant that remains platelet-associated.^{22–24} This plasma soluble form of GPVI (sGPVI) has been the marker for many quantitative studies on platelet activation through GPVI. The transmembrane metalloproteinases with protease activity, ADAM10 and ADAM17, have been identified as two sheddases that cleave the extracellular portion of GPVI independently of each other under various stimuli including the binding of GPVI ligands.²⁵ However, when antibody-mediated GPVI downregulation occurs, the catalyst for GPVI shedding appears to be independent of these MMPs, as antibody induced GPVI shedding occurred in mice depleted of ADAM10 and ADAM17.²⁵ Thus, it is likely that more sheddases participate in GPVI shedding and their role in stroke as a therapeutic target is yet to be determined.²⁶

GPVI and ischemic stroke

The importance of GPVI in hemostasis was first reported in a Japanese patient deficient in GPVI who had mild bleeding tendency and whose platelets failed to aggregate in response to collagen.²⁷ Since then it has

been found that platelets treated with GPVI-specific inhibitory antibodies show no interaction with collagen *in vivo*;²⁸ In addition, GPVI-deficient individuals exhibit a mild bleeding tendency²⁸ and GPVI-deficient mice show increased bleeding times.³⁰ These observations cement GPVI's role as the main signal generator leading to platelet activation, rather than functioning primarily as a platelet adhesion receptor,²⁹ but suggest that more work is needed to clarify the exact bleeding profile once GPVI is inhibited.

The exact role of GPVI in the different phenotypes of stroke is yet to be elucidated. We know that it plays a role in large artery atherosclerotic infarcts, but its role in lacunar and cardioembolic stroke is unclear. In large artery disease, elevated GPVI expression was shown to be associated with increased risk of stroke development. Enhanced GPVI expression is also seen after ischemic stroke and TIA, with these patients having a poorer clinical outcome at follow-up.³⁰ Elevated sGPVI levels were found in patients specifically after large artery infarcts and these levels decrease after 3–6 months, which may highlight a role for sGPVI measurements in this stroke substrate, but the evidence in cardioembolic and lacunar types were less convincing.³¹

The pathogenesis of lacunar infarcts is believed to be due to progressive ischemic leukoaraiosis caused by a genetic susceptibility to inflammation-mediated cerebrovascular injury in combination with the classic atherosclerotic risk factors.³² There is evidence that chronic endothelial dysfunction and activation leading to a prothrombotic environment may cause progression of leukoaraiosis. This is evidenced by higher levels of prothrombotic proteins such as ICAM1, thrombomodulin, fibrinogen, tissue factor in patients with cerebral leukoaraiosis compared to controls.^{33–35} Recently, thrombogenic fibrin was shown to be an activator of GPVI in mice.³⁶ Therefore, we could postulate that GPVI is involved in lacunar stroke and small vessel disease. Nevertheless, further studies are needed looking at GPVI in this subtype of stroke, where currently aspirin and clopidogrel play an important role in secondary prevention.

Recently, functional GPVI has been implicated as a possible receptor for polymerized fibrin, propagating thrombin generation.³⁷ Further studies may reveal the role of GPVI in cardioembolic stroke as well as thrombus propagation, clot stabilization and infarct growth after ischemic stroke.³⁸

Increased GPVI dimerization/multimerization could be one of the earliest measurable steps in platelet activation after plaque rupture. Studies are underway to measure GPVI dimer levels after acute stroke in comparison to healthy controls, which may be a useful diagnostic tool in the future. Since the dimer is the functional receptor form of GPVI, this may partially

explain why the studies discussed above measuring total GPVI (dimers plus monomers) did not find any correlation between its levels and severity of stroke.³⁰

Targeting GPVI in ischemic stroke: Beyond antiplatelet therapy

Current antiplatelet therapy acts on this cascade of platelet activation via different methods and is licenced for both acute coronary syndrome (ACS) and ischemic stroke. Aspirin irreversibly inhibits both COX 1 and 2, thus stopping TxA₂ formation after platelet activation. Clopidogrel and ticagrelor both irreversibly block the ADP receptor P2Y₁₂.³⁹ Thus both these classes of antiplatelet drugs work by inhibiting platelet aggregation. In comparison with single antiplatelet therapy, dual antiplatelets do convey a reduction in early stroke recurrence, combined transient ischemic attack (TIA), stroke and ACS, and all death⁴⁰ but dual antiplatelet therapy is reserved for high-risk individuals after TIA or stroke due to the risk of hemorrhage.

Studies on whether other platelet glycoprotein receptors could be targets for antithrombotic therapy have yielded less promising results. Abciximab, the Fab fragment of a chimeric mouse/human monoclonal antibody, is licenced in ACS for patients awaiting percutaneous coronary intervention. It antagonizes GPIIb/IIIa, which is the final mediator in the pathway to platelet aggregation. Unfortunately it has shown no benefit in functional outcome in acute stroke, coupled with

significant increases in fatal or symptomatic intracranial hemorrhage.⁴¹

Due to the fact that GPIb α (part of the GPIb-IX-V complex) plays a crucial role in platelet adhesion to endothelium in high shear conditions, it has become an attractive target for potential pharmaceutical development. GPIb α and GPVI are closely linked on the platelet surface and are thought to activate similar signaling processes.⁴² Much of the knowledge on platelet receptors and ischemic stroke has come from experimentation on animal models, particularly rats and mice, by causing transient middle cerebral artery occlusion (tMCAO). Blockade of GPIb α using Fab fragments, although reducing infarct volume after tMCAO, led to prolonged tail bleeding times compared to those treated with the monoclonal antibody (mAb) anti-GPVI JAQ1.⁴³

From similar studies in mice, GPVI was shown to have a huge role in the formation of arterial thrombi. After tMCAO, GPVI-depleted mice (via JAQ1) demonstrate significantly reduced brain infarct volumes, no hemorrhagic transformation, normal platelet counts, and only moderately increased tail bleeding times.^{28,43} As platelets are anucleate cells and cannot synthesis protein de novo, injection of rat anti-GPVI antibodies (JAQ 1, 2, and 3) into mice in other studies offered long-term depletion of GPVI from murine platelet surfaces.^{28,44} Furthermore, the Fab fragments of antibodies 5C4,⁴⁵ OM2,⁴⁶ OM4,⁴⁷ 9O12,⁴⁸ and mFab-F¹⁶ are also potent inhibitors of GPVI-mediated platelet activation (Table 1). The evidence for antibody-driven

Table 1. Antibodies that inhibit GPVI-mediated activation

	Type	Effects
JAQ 1 ²⁸	Rat anti-mouse GPVI	Abolished platelet response to collagen or CRP Significantly reduced infarct size in tMCAO mice <i>Thrombocytopenia and moderate bleeding-time prolongation</i>
9O12 (Fab) ⁴⁸	Mouse mAb against human GPVI	Protects against arterial thrombus propagation and collagen-induced aggregation in vitro and ex vivo <i>Mild bleeding-time prolongation</i>
5C4 (Fab) ⁴⁵	Rat mAb against human GPVI	Almost complete inhibition of platelet aggregation in vivo
OM2 (Fab) ⁴⁶	Mouse mAb against human GPVI	Inhibits collagen-induced platelet aggregation <i>Mild bleeding-time prolongation</i>
OM4 (Fab) ⁴⁷	Mouse mAb against human GPVI	Inhibits collagen-induced platelet aggregation in vitro and ex vivo Reduces in vivo thrombosis in rat carotid artery model <i>No bleeding-time prolongation</i>
10B12 ⁵⁸	Human scFv against human GPVI	Inhibits CRP- and collagen-induced platelet aggregation in vitro
m-Fab-F ¹⁶	Human Fab against human GPVI	GPVI dimer specific Inhibits aggregation and platelet adhesion on collagen

GPVI blockade as a target for stroke therapy is increasing, with further efforts being made to attempt clearer translation into clinical medicine. Recently, Kraft et al.⁴⁹ successfully showed that GPVI inhibition using JAQ 1 in adult mice with diabetes and hypertension resulted in smaller cerebral infarcts, better functional outcome, and reduced intracerebral hemorrhage rates compared to mice treated with anti-GPIIb/IIIa.⁴⁹

Patients with autoantibodies to GPVI-FcR γ chain, causing its clearance from the platelet surface are seen to have inefficiently formed thrombi on collagen-coated surfaces.⁵⁰ However, their clinical presentation of mild thrombocytopenia, ecchymosis, epistaxis, and prolonged bleeding times highlights the difficulty in translating experimental findings into clinical practice.

Building on the work initially done on animal models,⁴⁵ Revacept (PR-15, GPVI-Fc), a recombinant, soluble fusion protein between the human extracellular collagen-binding domain of GPVI and the C-terminal of human Fc, has been developed to competitively inhibit GPVI binding sites on exposed collagen. It was tested in a phase I trial, after intravenous administration to 30 healthy males, in 2011⁵¹ and showed clear dose-dependent inhibition of collagen responsiveness and platelet aggregation. This was done without any significant effect on hemostasis and no significant thrombocytopenia.

Revacept appears to avoid bleeding tendencies by blocking both the GPVI and VWF-mediated platelet activation processes specifically at the site of collagen exposure, thereby avoiding prolongation of bleeding times as described with antibodies to GPVI.^{28,43} However, this has only been confirmed on trials in mice after left MCA lesions.⁵² Revacept also seems to exert an anti-atherosclerotic effect via vessel endothelial remodelling and a role as a primary preventative medication in the absence of plaque rupture may emerge.⁵³ It is currently undergoing phase II trials on patients with symptomatic carotid artery stenosis.⁵⁴

Losartan, the angiotensin II receptor blocker has previously shown anti-atherosclerotic effects by blocking inflammogenic mediators and also anti-aggregatory effects independent of its effects on hypertension.⁵⁵ EXP3179, its active metabolite, selectively inhibits GPVI function and therefore platelet adhesion and aggregation both in vitro and in vivo.⁵⁶ This occurs when losartan binds to the IgG-like domain of GPVI⁵⁷ and this interaction, coupled with the fact that losartan has no effect on bleeding profiles, may be the basis of a therapeutic target in the future.

Conclusion

The platelet–collagen interaction, and subsequent platelet activation and thrombus formation, can lead to devastating neurological damage via ischemic

stroke. There is a growing body of evidence pointing towards the importance of platelet glycoprotein receptors in stroke due to their requisite role in initiating downstream signaling mechanisms leading to platelet activation and pathological thrombus generation. Current understanding of thrombus formation suggests that GPVI is one such crucial receptor and inhibition of its collagen-induced signaling would be a specific pharmacological target for ischemic stroke. However, most of the research has been carried out in murine experimental models and may not necessarily translate to bleeding-free therapy in humans. The current clinical trials that are underway on Revacept are promising and could lead the way to a better understanding of GPVI antagonism in humans. Smaller studies are being carried out looking at novel small molecule inhibitors of GPVI and if a successful entity is discovered, the next decade should see the development of efficacious and highly specific therapies in ischemic stroke that have a large safety margin.

Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: Drs Jung and Warburton are recipients of a Special Project grant (2014–2018) from the British Heart Foundation for the Glycoprotein VI in Ischemic Stroke study (GYPSIE) that is currently underway (SP/13/7/30575). Dr Warburton receives funding from the Cambridge Biomedical Research Centre award (NIHR).

References

1. Thrift AG, Cadilhac DA, Thayabaranathan T, et al. Global stroke statistics. *Int J Stroke* 2014; 9: 6–18.
2. Adams H, Bendixen B, Kappelle L, et al. Classification of subtype of acute ischemic stroke. *Stroke* 1993; 23: 35–41.
3. CAPRIE Steering Committee. A randomised, blinded, trial of clopidogrel versus aspirin in patients at risk of ischaemic events (CAPRIE). *Lancet* 1996; 348: 1329–1339.
4. Diener H-C, Bogousslavsky J, Brass LM, et al. Aspirin and clopidogrel compared with clopidogrel alone after recent ischaemic stroke or transient ischaemic attack in high-risk patients (MATCH): Randomised, double-blind, placebo-controlled trial. *Lancet* 2004; 364: 331–337.
5. Jackson SP, Nesbitt WS and Kulkarni S. Signaling events underlying thrombus formation. *J Thromb Haemost* 2003; 1: 1602–1612.
6. Koziak K, Sévigny J, Robson SC, Siegel JB and Kaczmarek E. Analysis of CD39/ATP diphosphohydrolase (ATPDase) expression in endothelial cells, platelets and leukocytes. *Thromb Haemost* 1999; 82: 1538–1544.

7. Farndale RW, Siljander PR, Onley DJ, Sundaresan P, Knight CG and Barnes MJ. Collagen-platelet interactions: Recognition and signalling. *Biochem Soc Symp* 2003; 94: 81–94.
8. Lecut C, Schoolmeester A, Kuijpers MJE, et al. Principal role of glycoprotein VI in $\alpha 2\beta 1$ and $\alpha_{IIb}\beta_3$ activation during collagen-induced thrombus formation. *Arterioscler Thromb Vasc Biol* 2004; 24: 1727–1733.
9. Dorsam RT and Kunapuli SP. Central role of the P2Y₁₂ receptor in platelet activation. *J Clin Invest* 2004; 113: 340–345.
10. Clemetson JM, Polgar J, Magnenat E, Wells TN and Clemetson KJ. The platelet collagen receptor glycoprotein VI is a member of the immunoglobulin superfamily closely related to Fc α R and the natural killer receptors. *J Biol Chem* 1999; 274: 29019–29024.
11. Moroi M and Jung SM. Platelet glycoprotein VI: Its structure and function. *Thromb Res* 2004; 114: 221–233.
12. Michelson A. *Platelets*. London: Elsevier/Academic Press, 2013, pp. 215–222.
13. Miura Y, Takahashi T, Jung SM and Moroi M. Analysis of the interaction of platelet collagen receptor glycoprotein VI (GPVI) with collagen: A dimeric form of GPVI, but not the monomeric form, shows affinity to fibrous collagen. *J Biol Chem* 2002; 277: 46197–46204.
14. Nieswandt B, Bergmeier W, Schulte V, Rackebrandt K, Gessner JE and Zirngibl H. Expression and function of the mouse collagen receptor glycoprotein VI is strictly dependent on its association with the FcR γ chain. *J Biol Chem* 2000; 275: 23998–24002.
15. Berlanga O, Bori-Sanz T, James JR, et al. Glycoprotein VI oligomerization in cell lines and platelets. *J Thromb Haemost* 2007; 5: 1026–1033.
16. Jung SM, Tsuji K and Moroi M. Glycoprotein (GP) VI dimer as a major collagen-binding site of native platelets: Direct evidence obtained with dimeric GPVI-specific Fabs. *J Thromb Haemost* 2009; 7: 1347–1355.
17. Jung SM, Moroi M, Soejima K, et al. Constitutive dimerization of glycoprotein VI (GPVI) in resting platelets is essential for binding to collagen and activation in flowing blood. *J Biol Chem* 2012; 287: 30000–30013.
18. Morton LF, Hargreaves PG, Farndale RW, Young RD and Barnes MJ. Integrin $\alpha 2\beta 1$ -independent activation of platelets by simple collagen-like peptides: Collagen tertiary (triple-helical) and quaternary (polymeric) structures are sufficient alone for $\alpha 2\beta 1$ -independent platelet reactivity. *Biochem J* 1995; 306: 337–344.
19. Smethurst P, Onley DJ, Jarvis GE, et al. Structural basis for the platelet-collagen interaction: The smallest motif within collagen that recognizes and activates platelet Glycoprotein VI contains two glycine-proline-hydroxyproline triplets. *J Biol Chem* 2007; 282: 1296–1304.
20. Horii K, Kahn ML and Herr AB. Structural basis for platelet collagen responses by the immune-type receptor glycoprotein VI. *Blood* 2006; 108: 936–942.
21. Loyau S, Dumont B, Ollivier V, et al. Platelet glycoprotein VI dimerization, an active process inducing receptor competence, is an indicator of platelet reactivity. *Arterioscler Thromb Vasc Biol* 2012; 32: 778–785.
22. Bergmeier W, Rabie T, Strehl A, et al. GPVI down-regulation in murine platelets through metalloproteinase-dependent shedding. *J Thromb Haemost* 2004; 91: 951–958.
23. Gardiner EE, Karunakaran D, Shen Y, Arthur JF, Andrews RK and Berndt MC. Controlled shedding of platelet glycoprotein (GP)VI and GPIb-IX-V by ADAM family metalloproteinases. *J Thromb Haemost* 2007; 5: 1530–1537.
24. Al-Tamimi M, Mu FT, Arthur JF, et al. Anti-glycoprotein VI monoclonal antibodies directly aggregate platelets independently of Fc γ RIIIa and induce GPVI ectodomain shedding. *Platelets* 2009; 20: 75–82.
25. Bender M, Hofmann S, Stegner D, et al. Differentially regulated GPVI ectodomain shedding by multiple platelet expressed proteinases. *Blood* 2010; 116: 3347–3355.
26. Newman PJ. GPVI and the not so eager cleaver. *Blood* 2010; 116: 3124–3126.
27. Moroi M, Jung SM, Okuma M and Shinmyozu K. A patient with platelets deficient in glycoprotein VI that lack both collagen-induced aggregation and adhesion. *J Clin Invest* 1989; 84: 1440–1445.
28. Nieswandt BB, Schulte V, Bergmeier W, et al. Long-term antithrombotic protection by in vivo depletion of platelet glycoprotein VI in mice. *J Exp Med* 2001; 193: 459–469.
29. Misra A and Dikshit M. Platelets and arterial thrombosis: Evolving role of platelet GPVI as an antithrombotic drug target. *Proc Indian Natl Sci Acad* 2014; 80: 289.
30. Bigalke B, Stellos K, Geisler T, et al. Expression of platelet glycoprotein VI is associated with transient ischemic attack and stroke. *Eur J Neurol* 2010; 17: 111–117.
31. Al-Tamimi M, Gardiner EE, Thom JY, et al. Soluble glycoprotein VI is raised in the plasma of patients with acute ischemic stroke. *Stroke* 2011; 42: 498–500.
32. Behrouz R, Malek AR and Torbey MT. Small vessel cerebrovascular disease: The past, present, and future. *Stroke Res Treat* 2012; 2012.
33. Hassan A, Hunt BJ, O’Sullivan M, et al. Markers of endothelial dysfunction in lacunar infarction and ischaemic leukoaraiosis. *Brain* 2003; 126: 424–432.
34. Knottnerus ILH, Govers-Riemslog JWP, Hamulyak K, et al. Endothelial activation in lacunar stroke subtypes. *Stroke* 2010; 41: 1617–1622.
35. van Oijen M, Wittman JC, Hofman A, Koudstaal PJ and Breteler MMB. Fibrinogen is associated with an increased risk of Alzheimer disease and vascular dementia. *Stroke* 2005; 36: 2637–2641.
36. Alshehri OM, Hughes CE, Montague S, et al. Fibrin activates GPVI in human and mouse platelets. *Blood* 2015; 126: 1601–1608.
37. Mammadova-Bach E, Ollivier V, Loyau S, et al. Platelet glycoprotein VI binds to polymerized fibrin and promotes thrombin generation. *Blood* 2015; 26: 683–689.
38. Nieswandt B, Pleines I and Bender M. Platelet adhesion and activation mechanisms in arterial thrombosis and ischaemic stroke. *J Thromb Haemost* 2011; 9: 92–104.
39. Maffrand JP. The story of clopidogrel and its predecessor, ticlopidine: Could these major antiplatelet and antithrombotic drugs be discovered and developed today? *Comptes Rendus Chim* 2012; 15: 737–743.

40. Geeganage CM, Diener H-C, Algra A, et al. Dual or mono antiplatelet therapy for patients with acute ischemic stroke or transient ischemic attack: Systematic review and meta-analysis of randomized controlled trials. *Stroke* 2012; 43: 1058–1066.
41. Adams HP, Effron MB, Torner J, et al. Emergency administration of abciximab for treatment of patients with acute ischemic stroke: Results of an international phase III trial: Abciximab in Emergency Treatment of Stroke Trial (AbESTT-II). *Stroke* 2008; 39: 87–99.
42. Gardiner EE. Targeting GPVI as a novel antithrombotic strategy. *J Blood Med* 2014; 5: 59–68.
43. Kleinschnitz C, Pozgajova M, Pham M, Bendszus M, Nieswandt B and Stoll G. Targeting platelets in acute experimental stroke: Impact of glycoprotein Ib, VI, and IIb/IIIa blockade on infarct size, functional outcome, and intracranial bleeding. *Circulation* 2007; 115: 2323–2330.
44. Massberg S, Gawaz M, Gruner S, et al. A crucial role of glycoprotein VI for platelet recruitment to the injured arterial wall in vivo. *J Exp Med* 2002; 197: 41–49.
45. Massberg S, Konrad I, Bültmann A, et al. Soluble glycoprotein VI dimer inhibits platelet adhesion and aggregation to the injured vessel wall in vivo. *FASEB J* 2004; 18: 397–399.
46. Matsumoto Y, Takizawa H, Gong X, et al. Highly potent anti-human GPVI monoclonal antibodies derived from GPVI knockout mouse immunization. *Thromb Res* 2007; 119: 319–329.
47. Li H, Lockyer S, Concepcion A, et al. The Fab fragment of a novel anti-GPVI monoclonal antibody, OM4, reduces in vivo thrombosis without bleeding risk in rats. *Arterioscler Thromb Vasc Biol* 2007; 27: 1199–1205.
48. Lecut C, Feeney LA, Kingsbury G, et al. Human platelet glycoprotein VI function is antagonized by monoclonal antibody-derived Fab fragments. *J Thromb Haemost* 2003; 1: 2653–2662.
49. Kraft P, Schuhmann MK, Fluri F, et al. Efficacy and safety of platelet glycoprotein receptor blockade in aged and comorbid mice with acute experimental stroke. *Stroke* 2015; 46: 3502–3506.
50. Boylan B, Chen H, Rathore V, et al. Anti-GPVI-associated ITP: An acquired platelet disorder caused by auto-antibody-mediated clearance of the GPVI/FcR γ -chain complex from the human platelet surface. *Blood* 2004; 104: 1350–1355.
51. Ungerer M, Rosport K, Bültmann A, et al. Novel anti-platelet drug Revacept (dimeric glycoprotein VI-Fc) specifically and efficiently inhibited collagen-induced platelet aggregation without affecting general hemostasis in humans. *Circulation* 2011; 123: 1891–1899.
52. Goebel S, Li Z, Vogelmann J, et al. The GPVI-Fc fusion protein revacept improves cerebral infarct volume and functional outcome in stroke. *PLoS One* 2013; 8: e66960.
53. Bültmann A, Li Z, Wagner S, et al. Impact of glycoprotein VI and platelet adhesion on atherosclerosis—a possible role of fibronectin. *J Mol Cell Cardiol* 2010; 49: 532–542.
54. Poppert H. Revacept in symptomatic carotid stenosis (Revacept/CS/02). <http://clinicaltrials.gov/show/NCT01645306>. NLM Identifier: NCT01645306 (accessed 31 May 2015).
55. Krämer C, Sunkomat J, Witte J, et al. Angiotensin II receptor-independent antiinflammatory and antiaggregatory properties of losartan: Role of the active metabolite EXP3179. *Circ Res* 2002; 90: 770–776.
56. Grothusen C, Umbreen S, Konrad I, et al. EXP3179 inhibits collagen-dependent platelet activation via glycoprotein receptor-VI independent of AT1-receptor antagonism: Potential impact on atherothrombosis. *Arterioscler Thromb Vasc Biol* 2007; 27: 1184–1190.
57. Ono K, Ueda H, Yoshizawa Y, et al. Structural basis for platelet antiaggregation by angiotensin II type 1 receptor antagonist losartan (DuP-753) via glycoprotein VI. *J Med Chem* 2010; 53: 2087–2093.
58. Smethurst P, Joutsu-Korhonen L, O'Connor MN, et al. Identification of the primary collagen-binding surface on human glycoprotein VI by site-directed mutagenesis and by a blocking phage antibody. *Blood* 2004; 103: 903–911.